

## Answers to Odd-Numbered Questions

### CHAPTER 1

#### Section 1.1

##### Quick Review 1.1

1. -2    3. -1    5. (a) Yes    (b) No



7.  $\sqrt{2}$     9.  $y = \frac{4}{3}x - \frac{7}{3}$

##### Exercises 1.1

1.  $\Delta x = -2, \Delta y = -3$     3.  $\Delta x = -5, \Delta y = 0$

5. (a) and (c), (b) 3

7. (a) and (c), (b) 0




9.  $x = 3; y = 2$     11.  $x = 0; y = -\sqrt{2}$     13.  $y = 1(x - 1) + 1$

15.  $y = 2(x - 0) + 3$     17.  $y = 3x - 2$


19.  $y = -\frac{1}{2}x - 3$     21.  $3x - 2y = 0$

23.  $x = -2$     25.  $y = \frac{5}{2}x$

27. (a)  $-\frac{3}{4}$     (b) 3



29. (a)  $-\frac{4}{3}$     (b) 4    (c)



31. (a)  $y = -x$     (b)  $y = x$     33. (a)  $x = -2$     (b)  $y = 4$

35.  $m = \frac{7}{2}, b = -\frac{3}{2}$     37.  $y = -1$

39.  $y = 1(x - 3) + 4$

$y = x - 3 + 4$


$y = x + 1$ , which is the same equation.

41. (a)  $k = 2$     (b)  $k = -2$

43. 5.97 atmospheres ( $k = 0.0994$ )

45. (a)  $y = 2216.2x - 4387470.6$

(b) 2216.2; it represents the approximate rate of increase in earnings in dollars per year.



[1995, 2005] by [40000, 50000]

(d) about \$62,659

47. False. A vertical line has no slope.

49. A    51. D




53. (a)  $y = 5980x - 11,810,220$

(b) The rate at which the median price is increasing in dollars per year

(c)  $y = 21650x - 43,105,030$

(d) South: \$5,980 per year, West: \$21,650 per year; more rapidly in the West

55. The coordinates of the three missing vertices are  $(5, 2)$ ,  $(-1, 4)$  and  $(-1, -2)$ .



57.  $y = -\frac{3}{4}(x - 3) + 4$  or  $y = -\frac{3}{4}x + \frac{25}{4}$

#### Section 1.2

##### Quick Review 1.2

1.  $[-2, \infty)$     3.  $[-1, 7]$     5.  $(-4, 4)$

7. Translate the graph of  $f$  2 units left and 3 units downward.

9. (a)  $x = -3, 3$     (b) No real solution

11. (a)  $x = 9$     (b)  $x = -6$

Exercises 1.2

1. (a)  $A(d) = \pi\left(\frac{d}{2}\right)^2$  (b)  $A(4) = 4\pi \text{ in}^2$   
 3. (a)  $S(e) = 6e^2$  (b)  $S(5) = 150 \text{ ft}^2$   
 5. (a)  $(-\infty, \infty); (-\infty, 4]$  (b)  $[1, \infty); [2, \infty)$
- 
- 
9. (a)  $(-\infty, 2) \cup (2, \infty); (-\infty, 0) \cup (0, \infty)$  (b)  $(-\infty, 0) \cup (0, \infty); (-\infty, 1) \cup (1, \infty)$
- 
- 
13. (a)  $(-\infty, \infty); (-\infty, \infty)$  (b)  $(-\infty, \infty); (-\infty, 1]$
- 
- 
17. (a)  $(-\infty, \infty); [0, \infty)$  (b)  $(-\infty, \infty); (-\infty, \infty)$
- 
- 
21. Even 23. Neither 25. Even 27. Odd 29. Neither
31.  $[-4.7, 4.7]$  by  $[-1, 6]$
- 
33.  $[-3.7, 5.7]$  by  $[-4, 9]$
- 
35. Because if the vertical line test holds, then for each x-coordinate, there is at most one y-coordinate giving a point on the curve. This y-coordinate would correspond to the value assigned to the x-coordinate. Since there's only one y-coordinate, the assignment would be unique.
37. No 39. Yes

41.  $f(x) = \begin{cases} x, & 0 \leq x \leq 1 \\ 2-x, & 1 < x \leq 2 \end{cases}$
43.  $f(x) = \begin{cases} 2-x, & 0 < x \leq 2 \\ \frac{5}{3} - \frac{x}{3}, & 2 < x \leq 5 \end{cases}$
45.  $f(x) = \begin{cases} -x, & -1 \leq x < 0 \\ 1, & 0 < x \leq 1 \\ \frac{3}{2} - \frac{x}{2}, & 1 < x < 3 \end{cases}$
47.  $f(x) = \begin{cases} 0, & 0 \leq x \leq \frac{T}{2} \\ \frac{2}{T}x - 1, & \frac{T}{2} < x \leq T \end{cases}$
49. (a)  $[-9.4, 9.4]$  by  $[-6.2, 6.2]$   
 (b) All reals (c)  $(-\infty, 2]$
51. (a)  $x^2 + 2$  (b)  $x^2 + 10x + 22$  (c) 2 (d) 22 (e) -2 (f)  $x + 10$
53. (a)  $g(x) = x^2$  (b)  $g(x) = \frac{1}{x-1}$  (c)  $f(x) = \frac{1}{x}$  (d)  $f(x) = x^2$
55. (a) Because the circumference of the original circle was  $8\pi$  and a piece of length  $x$  was removed. (b)  $r = \frac{8\pi - x}{2\pi} = 4 - \frac{x}{2\pi}$   
 (c)  $h = \sqrt{16 - r^2} = \frac{\sqrt{16\pi x - x^2}}{2\pi}$   
 (d)  $V = -\frac{1}{3}\pi r^2 h = \frac{(8\pi - x)^2 \sqrt{16\pi x - x^2}}{24\pi^2}$
57. False.  $f(-x) \neq f(x)$  59. B 61. D
63. (a) For  $f \circ g$ :  $[-10, 70]$  by  $[-10, 3]$   
 Domain:  $[0, \infty)$ ; Range:  $[-7, \infty)$   
 For  $g \circ f$ :  $[-3, 20]$  by  $[-4, 4]$   
 Domain:  $[7, \infty)$ ; Range:  $[0, \infty)$
- (b)  $(f \circ g)(x) = \sqrt{x} - 7$ ;  
 $(g \circ f)(x) = \sqrt{x - 7}$

Section 1.3

Quick Review 1.3

1. 2.924 3. 0.192  
 5. 1.8882 7. \$630.58  
 9.  $x^{-18}y^{-5} = \frac{1}{x^{18}y^5}$

Exercises 1.3


1.  $[-4, 4]$  by  $[-8, 6]$
- 
3.  $[-4, 4]$  by  $[-4, 8]$
- 
- Domain: All reals Range:  $(-\infty, 3)$  Domain: All reals Range:  $(-2, \infty)$
5.  $3^{4x}$  7.  $2^{-6x}$  9.  $\approx 2.322$  11.  $\approx -0.631$   
 13. (a) 15. (e) 17. (b)  
 19. (a) 1.032, 1.041, 1.034, 1.034, 1.029  
 (b) One possibility is  $2168(1.034)^u$ .  
 (c) 3348 thousand, or 3,348,000  
 21. After 19 years  
 23. (a)  $A(t) = 6.6\left(\frac{1}{2}\right)^{t/14}$   
 (b) About 38.1145 days later  
 25.  $\approx 11.433$  years 27.  $\approx 11.090$  years  
 29.  $\approx 19.108$  years 31.  $2^{48} \approx 2.815 \times 10^{14}$
33. 

x	y	$\Delta y$
1	-1	
2	1	2
3	3	2
4	5	2

 35. 

x	y	$\Delta y$
1	1	3
2	4	5
3	9	7
4	16	
37. Since  $\Delta x = 1$ , the corresponding value of  $\Delta y$  is equal to the slope of the line. If the changes in  $x$  are constant for a linear function, then the corresponding changes in  $y$  are constant as well.
39. (a)  $y = 14153.84(1.01963)^x$
- 
- $[-5, 25]$  by  $[-5000, 30000]$
- (b) Estimate: 22,133,000; the estimate exceeds the actual by 14,000.  
 (c)  $\approx 0.020$  or 2%  
 41. False. It is positive  $1/9$   
 43. D 45. B

65. (a) For  $f \circ g$ :  $[-10, 10]$  by  $[-10, 10]$   
 Domain:  $[-2, \infty)$ ; Range:  $[-3, \infty)$   
 For  $g \circ f$ :  $[-4.7, 4.7]$  by  $[-2, 4]$   
 Domain:  $(-\infty, -1] \cup [1, \infty)$ ; Range:  $[0, \infty)$
- (b)  $(f \circ g)(x) = (\sqrt{x+2})^2 - 3 = x - 1, x \geq -2$   
 $(g \circ f)(x) = \sqrt{x^2 - 1}$
67. (a)  $[-2, 2]$  by  $[-1.5, 1.5]$
- 
- (b)  $[-2, 2]$  by  $[-1.5, 1.5]$
- 
69. (a)  $[-2, 2]$  by  $[-1.3, 1.3]$
- 
- (b)  $[-2, 2]$  by  $[-1.3, 1.3]$
- 
71. (a)  $[-3, 3]$  by  $[-1, 3]$
- 
- (b) Domain of  $y_1$ :  $[0, \infty)$   
 Domain of  $y_2$ :  $(-\infty, 1]$   
 Domain of  $y_3$ :  $[0, 1]$
- (c) The results for  $y_1 - y_2$ ,  $y_2 - y_1$ , and  $y_1 \cdot y_2$  are the same as for  $y_1 + y_2$  above.  
 Domain of  $\frac{y_1}{y_2}$ :  $[0, 1)$  Domain of  $\frac{y_2}{y_1}$ :  $(0, 1]$
- (d) The domain of a sum, difference, or product of two functions is the intersection of their domains.  
 The domain of a quotient of two functions is the intersection of their domains with any zeros of the denominator removed.



[-5, 5] by [-2, 10]

In this window, it appears they cross twice, although a third crossing off-screen appears likely.

x	change in Y1	change in Y2
1		
2	3	2
3	5	4
4	7	8

(c)  $x = -0.7667, x = 2, x = 4$  (d)  $(-0.7667, 2) \cup (4, \infty)$

49.  $a = 0.5, k = 3$

### Quick Quiz (Sections 1.1–1.3)

1. C 3. E

### Section 1.4

#### Quick Review 1.4

1.  $y = -\frac{5}{3}x + \frac{29}{3}$  3.  $x = 2$

5. x-intercepts:  $x = -4$  and  $x = 4$  y-intercepts: none


7. (a) Yes (b) No (c) Yes

9. (a)  $t = \frac{-2x - 5}{3}$  (b)  $t = \frac{3y + 1}{2}$

#### Exercises 1.4

1. Graph (c). Window:  $[-4, 4]$  by  $[-3, 3], 0 \leq t \leq 2\pi$


3. Graph (d). Window:  $[-10, 10]$  by  $[-10, 10], 0 \leq t \leq 2\pi$



[-3, 3] by [-1, 3]

No initial or terminal point


(b)  $y = x^2$ ; all



[-3, 3] by [-2, 2]

Initial point: (1, 0)  
Terminal point: (-1, 0)


(b)  $x^2 + y^2 = 1$ ; upper half  
(or  $y = \sqrt{1 - x^2}$ ; all)



[-1, 5] by [-1, 3]

Initial point: (0, 0)  
Terminal point: None


(b)  $y = \sqrt{x}$ ; all (or  $x = y^2$ ; upper half)



[-3, 3] by [-2, 2]

Initial point: (-1, 0)  
Terminal point: (0, 1)


(b)  $x^2 + y^2 = 1$ ; upper half (or  
 $y = \sqrt{1 - x^2}$ ; all)



[-9, 9] by [-6, 6]

Initial and terminal point: (0, 5)


(b)  $y = 2x + 3$ ; all



[-6, 6] by [-2, 6]

Initial point: (4, 0)  
Terminal point: None


(b)  $y = -x + 4; x \leq 4$



[-4.7, 4.7] by [-3.1, 3.1]

Initial point: (0, 2)  
Terminal point: (0, -2)


(b)  $(\frac{x}{4})^2 + (\frac{y}{2})^2 = 1$ ; right half  
(or  $x = 2\sqrt{4 - y^2}$ ; all)



[-3, 3] by [-2, 2]

Initial point: (0, 1)  
Terminal point: (1, 0)

(b)  $y = -x + 1; (0, 1)$  to  $(1, 0)$



[-3, 3] by [-2, 2]

The curve is traced and retraced in both directions, and there is no initial or terminal point.

(b)  $y = -2x^2 + 1; -1 \leq x \leq 1$

23. Possible answer:  $x = -1 + 5t, y = -3 + 4t, 0 \leq t \leq 1$

25. Possible answer:  $x = t^2 + 1, y = t, t \leq 0$

27. Possible answer:  $x = 2 - 3t, y = 3 - 4t, t \geq 0$

29.  $1 < t < 3$  31.  $-5 \leq t < -3$

33. Possible answer:  $x = t, y = t^2 + 2t + 2, t > 0$

35. Possible answers:

(a)  $x = a \cos t, y = -a \sin t, 0 \leq t \leq 2\pi$

(b)  $x = a \cos t, y = a \sin t, 0 \leq t \leq 2\pi$

(c)  $x = a \cos t, y = -a \sin t, 0 \leq t \leq 4\pi$

(d)  $x = a \cos t, y = a \sin t, 0 \leq t \leq 4\pi$

37. False. It is an ellipse.

39. D 41. A

43. (a) The resulting graph appears to be the right half of a hyperbola in the first and fourth quadrants. The parameter  $a$  determines the  $x$ -intercept. The parameter  $b$  determines the shape of the hyperbola. If  $b$  is smaller, the graph has less steep slopes and appears "sharper." If  $b$  is larger, the slopes are steeper and the graph appears more "blunt."

(b) This appears to be the left half of the same hyperbola.

(c) Because both  $\sec t$  and  $\tan t$  are discontinuous at these points. This might cause the grapher to include extraneous lines (the asymptotes of the hyperbola) in its graph.

(d)  $(\frac{x}{a})^2 - (\frac{y}{b})^2 = (\sec t)^2 - (\tan t)^2 = 1$


by a standard trigonometric identity.

(e) This changes the orientation of the hyperbola. In this case,  $b$  determines the  $y$ -intercept of the hyperbola, and  $a$  determines the shape.

The parameter interval  $(-\frac{\pi}{2}, \frac{\pi}{2})$  gives the upper half of the

hyperbola. The parameter interval  $(\frac{\pi}{2}, \frac{3\pi}{2})$  gives the lower half.

The same values of  $t$  cause discontinuities and may add extraneous lines to the graph.



[-3, 3] by [-2, 2]

No initial or terminal point

(b)  $x^2 - y^2 = 1$ ; left branch  
(or  $x = -\sqrt{y^2 + 1}$ ; all)

47.  $x = 2 \cot t, y = 2 \sin^2 t, 0 < t < \pi$

### Section 1.5

#### Quick Review 1.5

1. 1 3.  $x^{2/3}$

5. Possible answer:  $x = t, y = \frac{1}{t-1}, t \geq 2$

7. (4, 5)

9. (a) (1.58, 3) (b) No intersection

#### Exercises 1.5

1. No 3. Yes 5. Yes 7. Yes 9. No 11. No

13.  $f^{-1}(x) = \frac{x-3}{2}$


15.  $f^{-1}(x) = (x+1)^{1/3}$  or  $\sqrt[3]{x+1}$

17.  $f^{-1}(x) = -x^{1/2}$  or  $-\sqrt{x}$


19.  $f^{-1}(x) = 2 - (-x)^{1/2}$  or  $2 - \sqrt{-x}$

21.  $f^{-1}(x) = \frac{1}{x^{1/2}}$  or  $\frac{1}{\sqrt{x}}$


23.  $f^{-1}(x) = \frac{1-3x}{x-2}$




[-6, 6] by [-4, 4]



[-4.5, 4.5] by [-3, 3]



[-4.5, 4.5] by [-3, 3]




[-3, 3] by [-2, 2]

33.  $t = \frac{\ln 2}{\ln 1.045} \approx 15.75$

35.  $x = \ln\left(\frac{3 \pm \sqrt{5}}{2}\right) \approx -0.96$  or  $0.96$

37.  $y = e^{2t+4}$



[-10, 5] by [-7, 3]

Domain:  $(-\infty, 3)$   
Range: all reals


43.  $f^{-1}(x) = \log_2\left(\frac{x}{100-x}\right)$

45. (a)  $f(f(x)) = \sqrt{1 - (f(x))^2}$   
 $= \sqrt{1 - (1 - x^2)}$   
 $= \sqrt{x^2}$   
 $= x, \text{ since } x \geq 0$

(b)  $f(f(x)) = f\left(\frac{1}{x}\right) = \frac{1}{1/x} = x$  for all  $x \neq 0$

47. About 14.936 years. (If the interest is only paid annually, it will take 15 years.)

49. (a)  $y = 1.758 + 1.076 \ln(x)$



[-2, 10] by [0, 6]

(b) 4 trillion cubic feet  
(c) Sometime during 2009

51. (a) Suppose that  $f(x_1) = f(x_2)$ . Then  $mx_1 + b = mx_2 + b$ , which gives  $x_1 = x_2$  since  $m \neq 0$ .

(b)  $f^{-1}(x) = \frac{x-b}{m}$ ; the slopes are reciprocals.

(c) They are also parallel lines with nonzero slope.

(d) They are also perpendicular lines with nonzero.

53. False. Consider  $f(x) = x^2, g(x) = \sqrt{x}$ . Notice that  $(f \circ g)(x) = x$  but  $f$  is not one-to-one.

55. A 57. B

59. If the graph of  $f(x)$  passes the horizontal line test, so will the graph of  $g(x) = -f(x)$  since it's the same graph reflected about the  $x$ -axis.

61. (a) Domain: All reals  
Range: If  $a > 0$ , then  $(d, \infty)$   
If  $a < 0$ , then  $(-\infty, d)$   
(b) Domain:  $(c, \infty)$   
Range: All reals

### Section 1.6

#### Quick Review 1.6

1.  $60^\circ$  3.  $-\frac{2\pi}{9}$  5.  $x \approx 0.6435, x \approx 2.4981$

7.  $x \approx 0.7854$  (or  $\frac{\pi}{4}$ ),  $x \approx 3.9270$  (or  $\frac{5\pi}{4}$ )



9. f(-x) = (-x)^3 - 3(-x) = -x^3 + 3x

= -(x^3 - 3x) = -f(x)

The graph is symmetric about the origin because if a point (a, b) is on the graph, then so is the point (-a, -b).

Exercises 1.6

1. 5π/4 3. 1/2 radian or ≈28.65° 5. Even 7. Odd


9. sin θ = 8/17, tan θ = -8/15, csc θ = 17/8,

sec θ = -17/15, cot θ = -15/8

11. (a) 2π/3

(b) x ≠ kπ/3, for integers k

(c) (-∞, -5] ∪ [1, ∞)




[-2π/3, 2π/3] by [-8, 8]

13. (a) π/3

(b) x ≠ kπ/6, for odd integers k

(c) All reals



[-π/2, π/2] by [-8, 8]

15. Possible answers are:

(a) [0, 4π] by [-3, 3]

(b) [0, 4π] by [-3, 3]


(c) [0, 2π] by [-3, 3]

17. (a) π (b) 1.5 (c) [-2π, 2π] by [-2, 2]

19. (a) π (b) 3 (c) [-2π, 2π] by [-4, 4]

21. (a) 6 (b) 4 (c) [-3, 3] by [-5, 5]

23. (a) y = 1.543 sin(2468.635x - 0.494) + 0.438



[0, 0.01] by [-2.5, 2.5]

(b) Frequency = 392.9, so it must be a "G."

25. The portion of the curve y = cos x between 0 ≤ x ≤ π passes the horizontal line test so it is one-to-one.



[-3, 3] by [-2, 4]

27. π/6 radian or 30° 29. ≈ -1.3734 radians or -78.6901°

31. x ≈ 1.190 and x ≈ 4.332 33. x = π/6 and x = 5π/6

35. x = 7π/6 + 2kπ and x = 11π/6 + 2kπ, k any integer

37. cos θ = 15/17 sin θ = 8/17 tan θ = 8/15

sec θ = 17/15 csc θ = 17/8 cot θ = 15/8

39. cos θ = -3/5 sin θ = 4/5 tan θ = -4/3

sec θ = -5/3 csc θ = 5/4 cot θ = -3/4

41. √72/11 ≈ 0.771

43. (a) 37 (b) 365 (c) 101 (d) 25

(e) f(x) = 37 sin[2π/365(x - 101)] + 25

45. (a) cot(-x) = cos(-x)/sin(-x) = cos(x)/-sin(x) = -cot(x)

(b) Assume that f is even and g is odd.

Then f(-x) = f(x) and g(-x) = -g(x) so f/g is odd.

The situation is similar for g/f.

47. Assume that f is even and g is odd.

Then f(-x)g(-x) = f(x)[-g(x)] = -f(x)g(x) so fg is odd.

49. (a) y = 3.0014 sin(0.9996x + 2.0012) + 2.9999

(b) y = 3 sin(x + 2) + 3

51. False. The amplitude is 1/2.

53. B 55. A

57. (a) √2 sin(ax + π/4) (b) See part (a).

(c) It works.

(d) sin(ax + π/4)


= sin(ax) \* 1/√2 + cos(ax) \* 1/√2

= 1/√2 (sin ax + cos ax)

So, sin(ax) + cos(ax) = √2 sin(ax + π/4).

59. Since sin(x) has period 2π, (sin(x + 2π))^3 = (sin(x))^3. This function has period 2π. A graph shows that no smaller number works for the period.

61. One possible graph:



[-π/60, π/60] by [-2, 2]

Quick Quiz (Sections 1.4-1.6)

1. C 3. E

Review Exercises

1. y = 3x - 9 2. y = -1/2x + 3/2 3. x = 0 4. y = -2x

5. y = 2 6. y = -2/5x + 21/5 7. y = -3x + 3 8. y = 2x - 5

9. y = -4/3x - 20/3 10. y = -5/3x - 19/3 11. y = 2/3x + 8/3

12. y = 5/3x - 5 13. y = -1/2x + 3 14. y = -2/7x - 6/7


15. Origin 16. y-axis 17. Neither 18. y-axis

19. Even 20. Odd 21. Even 22. Odd 23. Odd

24. Neither 25. Neither 26. Even

27. (a) Domain: all reals


(b) Range: [-2, ∞)



[-10, 10] by [-10, 10]

29. (a) Domain: [-4, 4]


(b) Range: [0, 4]



[-9.4, 9.4] by [-6.2, 6.2]

31. (a) Domain: all reals


(b) Range: (-3, ∞)



[-4, 4] by [-5, 15]

33. (a) Domain: all reals


(b) Range: [-3, 1]



[-π, π] by [-5, 5]

35. (a) Domain: (3, ∞)


(b) Range: all reals



[-3, 10] by [-4, 4]

37. (a) Domain: [-4, 4]


(b) Range: [0, 2]



[-6, 6] by [-3, 3]

28. (a) Domain: (-∞, 1]


(b) Range: [-2, ∞)



[-9.4, 9.4] by [-3, 3]

30. (a) Domain: all reals


(b) Range: (1, ∞)



[-6, 6] by [-4, 20]

32. (a) Domain: x ≠ kπ/4, for odd integers k


(b) Range: all reals



[-π/2, π/2] by [-8, 8]

34. (a) Domain: all reals


(b) Range: [0, ∞)



[-8, 8] by [-3, 3]

36. (a) Domain: all reals


(b) Range: all reals



[-10, 10] by [-4, 4]

38. (a) Domain: [-2, 2]

(b) Range: [-1, 1]



[-3, 3] by [-2, 2]

39. f(x) = { 1-x, 0 ≤ x < 1; 2-x, 1 ≤ x ≤ 2 }

40. f(x) = { 5x/2, 0 ≤ x < 2; -5/2x + 10, 2 ≤ x ≤ 4 }

41. (a) 1 (b) 1/√2.5 (= √2/5) (c) x, x ≠ 0 (d) 1/√(1+√(x+2)+2)

42. (a) 2 (b) 1 (c) x (d) √(x+1)+1

43. (a) (f ∘ g)(x) = -x, x ≥ -2

(g ∘ f)(x) = √(4-x^2)

(b) Domain (f ∘ g): [-2, ∞)

Domain (g ∘ f): [-2, 2]

(c) Range (f ∘ g): (-∞, 2]

Range (g ∘ f): [0, 2]

44. (a) (f ∘ g)(x) = √(1-x)

(g ∘ f)(x) = √(1-√x)


(b) Domain (f ∘ g): (-∞, 1]

Domain (g ∘ f): [0, 1]

(c) Range (f ∘ g): [0, ∞)

Range (g ∘ f): [0, 1]

45. (a)




[-6, 6] by [-4, 4]

Initial point: (5, 0)

Terminal point: (5, 0)

(b) (x/5)^2 + (y/2)^2 = 1; all

46. (a)




[-9, 9] by [-6, 6]

Initial point: (0, 4)

Terminal point: (0, -4)

(b) x^2 + y^2 = 16; left half

47. (a)




[-8, 8] by [-10, 20]

Initial point: (4, 15)

Terminal point: (-2, 3)

(b) y = 2x + 7; from (4, 15) to (-2, 3)

48. (a)



[-8, 8] by [-4, 6]

Initial point: None

Terminal point: (3, 0)

(b) y = √(6-2x); all


49. Possible answer: x = -2 + 6t, y = 5 - 2t, 0 ≤ t ≤ 1

50. Possible answer: x = -3 + 7t, y = -2 + t, -∞ < t < ∞

51. Possible answer: x = 2 - 3t, y = 5 - 5t, 0 ≤ t


52. Possible answer: x = t, y = t(t - 4), t ≤ 2

53. (a) f^-1(x) = (2-x)/3



[-6, 6] by [-4, 4]

54. (a) f^-1(x) = √(x-2)



[-6, 12] by [-4, 8]

55.  $\approx 0.6435$  radians or  $36.8699^\circ$

56.  $\approx -1.1607$  radians or  $-66.5014^\circ$

57.  $\cos \theta = \frac{3}{7}$   $\sin \theta = \frac{\sqrt{40}}{7}$   $\tan \theta = \frac{\sqrt{40}}{3}$


$\sec \theta = \frac{7}{3}$   $\csc \theta = \frac{7}{\sqrt{40}}$   $\cot \theta = \frac{3}{\sqrt{40}}$

58. (a)  $x \approx 3.3430$  and  $x \approx 6.0818$


(b)  $x \approx 3.3430 + 2k\pi$  and  $x \approx 6.0818 + 2k\pi$ ,  $k$  any integer

59.  $x = -5 \ln 4$


60. (a)




(b)




(c)




(d)



61. (a)




(b)



62. (a)  $V = 100,000 - 10,000x$ ,  $0 \leq x \leq 10$  (b) After 4.5 years

63. (a) 90 units (b)  $90 - 52 \ln 3 \approx 32.8722$  units

(c)



64. After  $\frac{\ln(10/3)}{\ln 1.08} \approx 15.6439$  years


(If the bank only pays interest at the end of the year, it will take 16 years.)

65. (a)  $N = 4 \cdot 2^t$  (b) 4 days; 64; 1 week; 512

(c) After  $\frac{\ln 500}{\ln 2} \approx 8.9658$  days, or after nearly 9 days

(d) Because it suggests the number of guppies will continue to double indefinitely and become arbitrarily large, which is impossible due to the finite size of the tank and the oxygen supply in the water.


66. (a)  $y = 72.695x - 143,940.564$



(b) 2103

(c) Slope = 72.695. It represents the number of doctoral degrees earned per year.

67. (a)  $y = 19,092(1.0025)^x$



(b) 19,526 thousand or 19,526,000

(c) 0.0025 or 0.25%.

68. (a)  $m = -1$  (b)  $y = -x - 1$  (c)  $y = x + 3$  (d) 2

69. (a)  $(2, \infty)$  (b)  $(-\infty, \infty)$  (c)  $x = 2 + e \approx 4.718$

(d)  $f^{-1}(x) = 2 + e^{1-x}$

(e)  $(f \circ g^{-1})(x) = f(f^{-1}(x)) = f(2 + e^{1-x}) = 1 - \ln(2 + e^{1-x} - 2)$   
 $= 1 - \ln(e^{1-x})$   
 $= 1 - (1 - x)$   
 $= x$

$(f^{-1} \circ g)(x) = f^{-1}(f(x)) = f^{-1}(1 - \ln(x - 2)) = 2 + e^{1 - \ln(x - 2)}$   
 $= 2 + e^{\ln(x - 2)}$   
 $= 2 + (x - 2)$   
 $= x$

70. (a)  $(-\infty, \infty)$  (b)  $[-2, 4]$  (c)  $\pi$  (d) Even (e)  $x \approx 2.526$

## CHAPTER 2

### Section 2.1

#### Quick Review 2.1

1. 0 3. 0 5.  $-4 < x < 4$

7.  $-1 < x < 5$  9.  $x = 6$

#### Exercises 2.1

1. 48 ft/sec 3. 96 ft/sec

5.  $2c^3 - 3c^2 + c - 1$

7.  $-\frac{3}{2}$  9. -15 11. 0 13. 4

15. (a)

$x$	-0.1	-0.01	-0.001	-0.0001
$f(x)$	1.566667	1.959697	1.995997	1.999600

(b)

$x$	0.1	0.01	0.001	0.0001
$f(x)$	2.372727	2.039703	2.003997	2.000400

The limit appears to be 2.

17. (a)

$x$	-0.1	-0.01	-0.001	-0.0001
$f(x)$	-0.054402	-0.005064	-0.000827	-0.000031

(b)

$x$	0.1	0.01	0.001	0.0001
$f(x)$	-0.054402	-0.005064	-0.000827	-0.000031

The limit appears to be 0.

19. (a)

$x$	-0.1	-0.01	-0.001	-0.0001
$f(x)$	2.0567	2.2763	2.2999	2.3023

(b)

$x$	0.1	0.01	0.001	0.0001
$f(x)$	2.5893	2.3293	2.3052	2.3029

The limit appears to be approximately 2.3.

21. Expression not defined at  $x = -2$ . There is no limit.

23. Expression not defined at  $x = 0$ . There is no limit.

25.  $\frac{1}{2}$  27.  $-\frac{1}{2}$  29. 12 31. -1 33. 0

35. Answers will vary. One possible graph is given by the window  $[-4.7, 4.7]$  by  $[-15, 15]$  with  $X_{\text{sc1}} = 1$  and  $Y_{\text{sc1}} = 5$ .

37. 0 39. 0 41. 1

43. (a) True (b) True

(c) False (d) True

(e) True (f) True

(g) False (h) False

(i) False (j) False

45. (a) 3 (b) -2

(c) No limit (d) 1

47. (a) -4 (b) -4

(c) -4 (d) -4

49. (a) 4 (b) -3


(c) No limit (d) 4

51. (c) 53. (d)

55. (a) 6 (b) 0

(c) 9 (d) -3


57. (a)



$[-3, 6]$  by  $[-1, 5]$

(b) Right-hand: 2 Left-hand: 1  
 (c) No, because the two one-sided limits are different


59. (a)



$[-5, 5]$  by  $[-4, 8]$

(b) Right-hand: 4  
 Left-hand: no limit  
 (c) No, because the left-hand limit doesn't exist


61. (a)



$[-2\pi, 2\pi]$  by  $[-2, 2]$

(b)  $(-2\pi, 0) \cup (0, 2\pi)$   
 (c)  $c = 2\pi$  (d)  $c = -2\pi$

63. (a)



$[-2, 4]$  by  $[-1, 3]$

(b)  $(0, 1) \cup (1, 2)$  (c)  $c = 2$   
 (d)  $c = 0$

65. 0

67. 0

69. (a) 14.7 m/sec

(b) 29.4 m/sec

71. True. Definition of limit.

73. C 75. E

77. (a) Because the right-hand limit at zero depends only on the values of the function for positive  $x$ -values near zero

(b) Use: area of triangle =  $\left(\frac{1}{2}\right)$ (base)(height)

area of circular sector =  $\frac{(\text{angle})(\text{radius})^2}{2}$

(c) This is how the areas of the three regions compare.

(d) Multiply by 2 and divide by  $\sin \theta$ .

(e) Take reciprocals, remembering that all of the values involved are positive.

(f) The limits for  $\cos \theta$  and 1 are both equal to 1. Since  $\frac{\sin \theta}{\theta}$  is between them, it must also have a limit of 1.