

$\chi^2$ -w.s. (O.E.)

5. A rural county hospital offers several health services. The hospital administrators conducted a poll to determine whether the residents' satisfaction with the available services depends on their gender. A random sample of 1,000 adult county residents was selected. The gender of each respondent was recorded and each was asked whether he or she was satisfied with the services offered by the hospital. The resulting data are shown in the table below.

	Male	Female	Total
Satisfied	384	416	800
Not Satisfied	80	120	200
Total	464	536	1,000

- (a) Using a significance level of 0.05, conduct an appropriate test to determine if, for adult residents of this county, there is an association between gender and whether or not they were satisfied with services offered by the hospital.
- (b) Is  $\frac{800}{1,000}$  a reasonable estimate for the proportion of all adult county residents who are satisfied with the services offered by this hospital? Explain why or why not.

2. The Colorado Rocky Mountain Rescue Service wishes to study the behavior of lost hikers. If more were known about the direction in which lost hikers tend to walk, then more effective search strategies could be devised. Two hundred hikers selected at random from those applying for hiking permits are asked whether they would head uphill, downhill, or remain in the same place if they became lost while hiking. Each hiker in the sample was also classified according to whether he or she was an experienced or novice hiker. The resulting data are summarized in the following table.

	Direction		
	Uphill	Downhill	Remain in Same Place
Novice	20	50	50
Experienced	10	30	40

Do these data provide convincing evidence of an association between the level of hiking expertise and the direction the hiker would head if lost?

Give appropriate statistical evidence to support your conclusion.